

Applications of Four-Vectors in Particle Physics

By Tianze Zhang

AUTHOR BIO

Tianze Zhang (Chris) is a high school student in New Zealand with a strong interest in physics and mathematical reasoning. He enjoys using computational modelling to explore theoretical ideas and hopes to continue studying physics through an interdisciplinary lens.

ABSTRACT

Special Relativity requires a mathematical framework that treats space and time as components of a single four-dimensional structure. This paper examines the four-vector formalism as that framework, tracing its use from the derivation of the Lorentz transformations through applications in modern particle physics. This paper begins with the failure of Galilean relativity to accommodate Maxwell's prediction of a constant speed of light, and derives the Lorentz transformations that resolve this conflict. Time dilation is illustrated using the atmospheric muon experiment, which confirms the effect experimentally. Four-vectors are then introduced as quantities whose invariant length under Lorentz transformations makes them suited to describing energy, momentum, and spacetime intervals across reference frames. Using this formalism, this paper analyzes Compton scattering, showing how conservation of four-momentum produces the scattering formula and supports the particle interpretation of light. The paper extends to Feynman diagrams and the Mandelstam variables s , t , and u , which provide a Lorentz-invariant description of scattering processes used throughout collider physics. Each application illustrates the same underlying principle: four-vector algebra converts calculations that would otherwise depend on the observer's frame into results that hold universally. The paper closes by connecting this formalism to its role in contemporary high-energy physics, where scattering amplitudes and cross sections are computed directly in terms of invariant four-vector quantities.

Keywords: special relativity, four-vectors, Lorentz transformations, particle physics, Compton scattering, Mandelstam variables

I. INTRODUCTION

The history of physics is one of unification. Where Isaac Newton unified the falling apple with the orbiting moon, the early 20th century necessitated a reconciliation between the mechanics of moving bodies and the electrodynamics of fields. This report provides an analysis of the mathematical and physical framework that emerged from this reconciliation: the four-vector formalism of Special Relativity. We explore how the transition from a three-dimensional Euclidean worldview to a four-dimensional Minkowski spacetime is not merely a mathematical convenience but a fundamental requirement for describing the high-energy universe.

A. The Classical Foundation and Its Cracks

Classical mechanics, governed by Newton's laws, relies on the concept of absolute time. In the Newtonian paradigm, time is a universal parameter, t , flowing uniformly and independently of any observer. Space is a rigid, Euclidean container. The transformation between two inertial reference frames, frames moving at constant velocity relative to one another, is described by the Galilean transformation. If a frame S' moves with constant velocity v along the x -axis relative to frame S , the coordinates transform as:

$$x' = x - vt, \quad t' = t$$

This transformation implies that velocities add linearly: $u' = u - v$. For centuries, this relation held, accurately describing everything from ballistics to planetary orbits. However, the edifice of classical mechanics began to fracture with the advent of James Clerk Maxwell's equations of electromagnetism. Maxwell's theory predicted the existence of electromagnetic waves propagating at a fixed speed, c , roughly 3×10^8 m/s. A critical contradiction arose: if the Galilean velocity addition law were true, an observer chasing a light beam at speed v should measure the light's speed as $c - v$. Yet, Maxwell's equations contain no reference to the velocity of the source or the observer; c appears as a fundamental constant of nature, determined solely by the permittivity and permeability of the vacuum (ϵ_0, μ_0). This conflict suggested two possibilities: either Maxwell's equations were incomplete (requiring a medium to define a preferred frame), or the Galilean transformation was only an approximation valid at low velocities (DiSalle, 2020).

B. The Breakdown at High Velocities

The resolution, formalized by Albert Einstein in 1905, was to accept the constancy of the speed of light as a postulate (Einstein, 1905). This acceptance forces the abandonment of absolute time. If c is invariant, temporal measurements must vary between observers to compensate for spatial motion. Newtonian mechanics works exceptionally well as an approximation when velocities are small compared to light ($v \ll c$). In this limit, the Lorentz factor $\gamma \approx 1$, and relativistic equations reduce to classical ones. However, as $v \rightarrow c$, Newtonian mechanics fails catastrophically (Rivadulla, 2004; Tasson, 2019):

- **Velocity Addition:** Galilean addition allows velocities to exceed c , violating causality.

- **Inertia:** In classical mechanics, a constant force produces constant acceleration ($F = ma$). In relativity, as a particle approaches c , its resistance to acceleration increases effectively to infinity, preventing it from crossing the light barrier.
- **Energy Divergence:** The classical kinetic energy $K = \frac{1}{2}mv^2$ remains finite at any velocity. The relativistic energy diverges as $v \rightarrow c$, requiring infinite energy to accelerate a massive particle to light speed.

The four-vector formalism ensures there is no failure mathematically. Creating a single vector that has space in three dimensions and a fourth of time, this dictates that the equations must maintain their form. Maintaining their form simply means that the system being described looks the same in all inertial reference frames.

II. LORENTZ TRANSFORMATIONS AND CONSEQUENCES

The invariance of the speed of light requires a new set of transformations between inertial frames, replacing the Galilean relations. These transformations must preserve the causal structure of spacetime. That is, the geometric arena in which events are located by specifying both their spatial position and the time at which they occur to ensure that all inertial observers measure the same value of c . This leads naturally to the Lorentz transformations (Taylor & Wheeler, 1992).

A. Inertial Frames And The Standard Configuration

We consider two inertial frames S and S' in the standard configuration. Frame S' moves with constant velocity v along the x -axis of S , and the origins coincide at $t = t' = 0$. The coordinates of an event are denoted by (t, x, y, z) in S and (t', x', y', z') in S' .

B. Lorentz Transformations in 1+1 Dimensions

While the Galilean transformations $x' = x - vt$, $t' = t$, fail to preserve the speed of light, Einstein's postulates require that the speed of a light pulse be c in both S and S' . A light pulse emitted from the common origin at $t = t' = 0$ satisfies $c^2 t^2 - x^2 = 0$ in S and $c^2 t'^2 - x'^2 = 0$ in S' . More generally, if we demand that the transformations between inertial frames be linear (as required by the homogeneity of spacetime), then the quantity $s^2 = c^2 t^2 - x^2$ must be invariant under these transformations. This requirement, together with the assumption of linearity, uniquely determines the Lorentz transformations. For motion along the x -direction, they are given by

$$t' = \gamma \left(t - \frac{v}{c^2} x \right), x' = \gamma (x - vt), \text{ with the Lorentz factor } \gamma = \frac{1}{\sqrt{1 - v^2/c^2}}.$$

The transverse coordinates remain unchanged: $y' = y$, $z' = z$. It is straightforward to verify that these relations satisfy $c^2 t'^2 - x'^2 = c^2 t^2 - x^2$, illustrating that Lorentz transformations preserve the spacetime interval.

C. Matrix Form of Lorentz Transformations

It is convenient to write spacetime coordinates as a column vector, $X = (ct \ x \ y \ z)$, $X' = (ct' \ x' \ y' \ z')$. Then a Lorentz transformation can be expressed compactly as $X' = \Lambda X$, where Λ is the 4×4 Lorentz transformation matrix. For a boost along the x -axis, we have $\Lambda = (\gamma \ -\ \beta\gamma \ 0 \ 0 \ -\ \beta\gamma \ \gamma \ 0 \ 0 \ 0 \ 0 \ 0 \ 0 \ 1 \ 0 \ 0 \ 0 \ 1)$, $\beta = \frac{v}{c}$. These matrices satisfy the defining relation of the Lorentz group, $\Lambda^T \eta \Lambda = \eta$, where $\eta = \text{diag}(1, -1, -1, -1)$ is the Minkowski metric. This expresses the invariance of the interval in matrix language. In the following section, we will introduce index notation and the concept of four-vectors, which generalize this matrix structure to all physical quantities.

D. Simple Consequences of Lorentz Transformations

1. Time Dilation

Consider a clock at rest in frame S' , so that $\Delta x' = 0$ and the proper time interval is $\Delta \tau = \Delta t'$. Using the Lorentz transformation, $\Delta t = \gamma \Delta \tau$, we see that moving clocks run more slowly when observed from another inertial frame. This effect is essential in particle physics. A striking example is the cosmic-ray muon. Muons produced by cosmic-ray interactions in the upper atmosphere have a rest-frame half-life of approximately $1.56 \mu\text{s}$ (Navas et al., 2024). Without time dilation, a muon traveling at close to c would traverse only a few hundred meters before decaying, far less than the roughly 15 km from the upper atmosphere to the Earth's surface. However, because such muons are highly relativistic ($\gamma \sim 10\text{--}100$), their observed lifetime in the laboratory frame is dilated by the factor γ , allowing a significant fraction to reach sea level. This was among the earliest experimental confirmations of Special Relativity (Rossi & Hall, 1941).

2. Length Contraction

A rod of rest length L_0 (the length measured by an observer for whom the rod is stationary) is at rest in frame S . To measure its length in S' , the positions of its endpoints must be recorded simultaneously in S' , i.e., $\Delta t' = 0$. Applying the Lorentz transformation gives $L = \frac{L_0}{\gamma}$, showing that moving objects appear shortened along the direction of motion. Note that this contraction is intimately connected to the relativity of simultaneity: the requirement that the endpoints be measured at the same time in S' is essential, since observers in different frames disagree on what constitutes “the same time.” Length contraction is therefore not a mechanical compression of the object, but a consequence of the structure of spacetime itself.

3. Relativity of Simultaneity

Events that are simultaneous in one inertial frame don't have to be simultaneous in another; this holds true unless the events happen at the exact same place. We can see this in the equation below, if Δx is zero, then there would be no change in time. However, if the events do not take place at the exact location, then the two events could happen at another time in a different frame.

$$\Delta t' = -\gamma \frac{v}{c^2} \Delta x. \quad (1)$$

Thus, simultaneity is frame-dependent. We can see that time is dependent on location in the above equation. Therefore, there is a connection between space and time under Lorentz transformations, this is why the separation of space and time in Newtonian physics isn't applicable in all applications.

4. Relativistic Velocity Addition

Suppose a particle moves in frame S with velocity u_x , while frame S' moves with speed v relative to S . Differentiating the Lorentz transformation equations yields the velocity addition law,

$$u'_x = \frac{u_x - v}{1 - \frac{u_x v}{c^2}}.$$

This ensures that no matter how velocities combine, the resulting speed never exceeds c . For example, if both $u_x = 0.9c$ and $v = 0.9c$, Galilean addition would predict $u' = 1.8c$, while the relativistic formula gives $u' \approx 0.994c$, safely below the speed of light. In collider physics, where particles are viewed from different boosted frames, this formula is frequently essential.

5. Rapidity and Boost Composition

For high-energy physics it is often more convenient mathematically to parameterize boosts in terms of the rapidity θ , defined by

$$\tanh \theta = \beta, \quad \gamma = \cosh \theta, \quad \gamma \beta = \sinh \theta.$$

In this language, a boost along x axis takes the form

$$\Lambda(\theta) = \begin{pmatrix} \cosh \theta & -\sinh \theta & 0 & 0 \\ -\sinh \theta & \cosh \theta & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{pmatrix}$$

This matrix only includes the x axis and time, so there is no rotation or mathematical elements of two axis, such as x - y or z - x .

A significant advantage is that successive boosts along the same axis simply add:

$$\theta_{total} = \theta_1 + \theta_2.$$

This makes rapidity an effective tool in motion analysis and collisions.

Lorentz transformations as discussed above are critical when analyzing spacetime intervals. This is the rational to have these physical quantities that transform linearly and have invariant magnitudes because they are unchanged by the observer. Next, we will cover four vectors, the Minkowski metric and invariant quantities that fundamentally help define relativistic particle dynamics and scatter theory.

III. FOUR-VECTORS AND LORENTZ INVARIANTS

The defining feature of the Lorentz transformations is the preservation of the spacetime interval. This suggests that physical laws should be formulated in terms of quantities that transform linearly under Lorentz transformations and possess invariant scalar magnitudes. Such quantities are known as four-vectors. In relativistic particle physics, four-vectors provide the natural language for describing spacetime events, energy, momentum, and interactions in a manner that is independent of the observer's inertial frame (Griffiths, 2008; Halzen & Martin, 1984).

A. Four-Vectors and Index Notation

A four-vector A^μ is defined as an object whose components transform under a Lorentz transformation according to

$$A'^\mu = \Lambda^\mu_\nu A^\nu,$$

where Λ^μ_ν is a Lorentz transformation matrix. Throughout this report, Greek indices $\mu, \nu = 0, 1, 2, 3$ label spacetime components, while Latin indices $i, j = 1, 2, 3$ denote purely spatial components. Repeated indices are implicitly summed over, following the Einstein summation convention.

The geometry of spacetime is encoded in the Minkowski metric,

$$\eta_{\mu\nu} = \text{diag}(1, -1, -1, -1),$$

which allows indices to be raised and lowered:

$$A_\mu = \eta_{\mu\nu} A^\nu, \quad A^\mu = \eta^{\mu\nu} A_\nu.$$

Quantities constructed by contracting indices, such as $A^\mu B_\mu$, are Lorentz scalars and therefore invariant under changes of inertial frame.

B. The Spacetime Four-Vector X^μ

The position of an event in spacetime is described by the spacetime four-vector

$$X^\mu = (ct, x, y, z),$$

or, in covariant form,

$$X_\mu = (ct, -x, -y, -z).$$

Under a Lorentz transformation, X^μ transforms linearly, preserving the invariant spacetime interval $\Delta s^2 = \Delta X^\mu \Delta X_\mu = c^2 \Delta t^2 - \Delta x^2$.

The sign of Δs^2 classifies the separation between two events into three physically distinct categories.

If $\Delta s^2 > 0$, the separation is timelike. In this case there exists an inertial frame in which the two events occur at the same spatial position but at different times. Timelike-separated events can be causally connected by the motion of a massive particle, and the proper time between them is real and nonzero.

If $\Delta s^2 < 0$, the separation is spacelike. There exists a frame in which the events are simultaneous but occur at different spatial locations. No causal influence can propagate between spacelike-separated events without exceeding the speed of light, and the temporal order of such events is frame-dependent.

If $\Delta s^2 = 0$, the separation is lightlike or null. The events lie on the light cone and can be connected only by signals traveling at the speed of light. Lightlike intervals define the causal boundary between timelike and spacelike regions and play a fundamental role in relativistic field theory.

C. Proper Time and Four-Velocity

For a particle moving along a worldline $X^\mu(\tau)$, the invariant interval defines the proper time τ via

$$c^2 d\tau^2 = c^2 dt^2 - dx^2.$$

Proper time is the time measured in the instantaneous rest frame of the particle and is a Lorentz-invariant quantity.

The four-velocity is defined as the derivative of the spacetime four-vector with respect to proper time:

$$U^\mu = \frac{dX^\mu}{d\tau}.$$

In terms of the three-velocity v , this takes the explicit form

$$U^\mu = \gamma(c, v),$$

where $\gamma = (1 - v^2/c^2)^{-1/2}$. A key property of the four-velocity is that its Minkowski norm is invariant:

$$U^\mu U_\mu = c^2.$$

This result holds for all massive particles, regardless of their state of motion, and highlights the geometric nature of relativistic dynamics.

D. Four-Momentum P^μ

In relativistic mechanics, energy and momentum are unified into the four-momentum,

$$P^\mu = mU^\mu = \left(\frac{E}{c}, p\right),$$

where m is the invariant (rest) mass of the particle, E is its total energy, and p is its three-momentum. The covariant form is

$$P_{\mu} = \left(\frac{E}{c}, -p \right).$$

From the definition $P^{\mu} = mU^{\mu} = m\gamma(c, v)$, we can read off the explicit expressions

$$E = \gamma mc^2, \quad p = \gamma mv.$$

In the non-relativistic limit ($v \ll c$), the Lorentz factor admits a Taylor expansion $\gamma \approx 1 + \frac{1}{2} \frac{v^2}{c^2} + \dots$, so that

$$E = \gamma mc^2 \approx mc^2 + \frac{1}{2} mv^2 + \dots,$$

recovering the rest energy plus the classical kinetic energy.

E. Energy–Momentum Invariant

The contraction of the four-momentum with itself yields a Lorentz-invariant scalar:

$$P^{\mu} P_{\mu} = \frac{E^2}{c^2} - p^2.$$

Evaluating this quantity in the particle's rest frame, where $p = 0$ and $E = mc^2$, gives

$$P^{\mu} P_{\mu} = m^2 c^2.$$

Since this quantity is invariant, it must take the same value in all inertial frames. This leads directly to the relativistic energy–momentum relation,

$$E^2 = p^2 c^2 + m^2 c^4.$$

For massless particles such as photons ($m = 0$), this reduces to $E = pc$, demonstrating that the four-vector formalism naturally accommodates both massive and massless particles.

F. Role of Invariants in Particle Physics

Lorentz invariants play a central role in particle physics because they allow physical processes to be analyzed independently of the observer's frame. The invariant mass of a particle or system can be determined from measurements of energy and momentum in any frame. Furthermore, four-momentum conservation,

$$\sum_{\text{initial}} P^{\mu} = \sum_{\text{final}} P^{\mu},$$

provides a compact and frame-independent statement of the conservation laws governing particle interactions. Threshold energies, scattering kinematics, and decay products are most easily understood by evaluating Lorentz invariants to simplify the mathematics. Then once solving the calculations in a simpler frame, one can convert the data back to any other frame. In the following section, we apply the four-vector formalism to particle interactions, including relativistic collisions, photon–particle scattering, and the kinematic structure underlying Feynman diagrams.

IV. PARTICLE INTERACTIONS AND KINEMATICS

The power of the four-vector formalism becomes most apparent when analyzing particle interactions. In classical mechanics, energy and momentum are conserved separately. In relativistic physics, these are unified into the conservation of four-momentum. For any isolated system of particles, the total four-momentum is conserved:

$$\sum_{initial} P_i^\mu = \sum_{final} P_f^\mu.$$

This single vector equation encompasses both the conservation of relativistic energy (the temporal component, $\mu = 0$) and the conservation of three-momentum (the spatial components, $\mu = 1, 2, 3$).

The utility of this formalism lies in the Lorentz invariance of the scalar product $P^\mu P_\mu$. Because this quantity is frame-independent, we can evaluate the kinematics of a collision in the frame where the algebra is simplest, typically the Center of Momentum (CM) frame, and the results will hold true for an observer in the laboratory frame.

A. Colliders vs. Fixed Target Experiments

One of the most consequential applications of relativistic kinematics is the comparison between fixed-target experiments and colliding-beam experiments. Consider the goal of creating a heavy particle of mass M from the collision of two lighter particles of mass m .

1. Fixed Target Scenario

Consider a beam particle with energy E_{lab} and mass m colliding with a stationary target particle of mass m . The four-momenta in the laboratory frame are:

$$P_{beam} = \left(\frac{E_{lab}}{c}, p_{lab} \right),$$
$$P_{target} = (mc, 0).$$

The square of the total four-momentum is an invariant whose value equals E_{cm}^2/c^2 , where E_{cm} is the total energy in the CM frame (we will formalize this as the Mandelstam variable s in Section V B):

$$\frac{E_{cm}^2}{c^2} = (P_{beam} + P_{target})^2 = P_{beam}^2 + P_{target}^2 + 2P_{beam} \cdot P_{target}.$$

Using the invariant mass relation $P^2 = m^2 c^2$:

$$\frac{E_{cm}^2}{c^2} = m^2 c^2 + m^2 c^2 + 2 \left(\frac{E_{lab}}{c} \cdot mc - p_{lab} \cdot 0 \right).$$

Simplifying, we find:

$$E_{cm}^2 = 2m^2 c^4 + 2mc^2 E_{lab}.$$

In the high-energy limit where $E_{lab} \gg mc^2$, the available energy for particle creation scales as the square root of the beam energy:

$$E_{cm} \approx \sqrt{2mc^2 E_{lab}}.$$

This is highly inefficient. To double the available energy, one must increase the beam energy by a factor of four.

2. Collider Scenario

Now consider two particles approaching each other with equal and opposite momenta, effectively observing the collision from the center of momentum frame.

$$P_1 = \left(\frac{E}{c}, p \right),$$
$$P_2 = \left(\frac{E}{c}, -p \right).$$

The total four-momentum is simply $P_{tot} = (2E/c, 0)$. The available energy is:

$$E_{cm} = 2E.$$

These equations show particles one and two traveling at the same momentum but in opposite directions, thus when they become one new system upon collision the energy doubles on the new system and the particles are no longer moving in space with a net momentum of zero. Thus, there is no loss of energy from a momentum change.

In a collider, the available energy scales linearly with the beam energy. This kinematic advantage is the primary motivation for building colliders like the LHC; a “head-on” collision utilizes all the kinetic energy for new physics, whereas a fixed-target collision “wastes” a vast amount of energy primarily to conserve the total momentum of the system.

B. Threshold Energies

A common problem in particle physics is calculating the minimum (threshold) beam energy required to produce a particular final state. Consider the interaction:

$$p + p \rightarrow p + p + \pi^0.$$

What is the minimum energy with which a proton needs to strike a second, stationary proton, in order to create a neutral pion?

The threshold condition is defined in the CM frame: the minimum energy configuration occurs when all final particles are created at rest relative to each other. Thus, the final state effectively moves as a single object with mass equal to the sum of the product masses, $M_{final} = 2m_p + m_{\pi}$.

We equate the invariant $(P_{total})^2$ calculated in the initial Lab frame to the invariant in the final threshold configuration:

$$(P_{total})_{initial}^2 = (P_{total})_{final}^2$$

From the fixed target calculation above, $(P_{total})_{initial}^2 = 2m_p^2 c^2 + 2m_p E_{lab}$. At threshold, the final four-momentum is just the rest energy of the product masses squared:

$$(P_{total})_{final}^2 = (2m_p + m_\pi)^2 c^2.$$

Equating them:

$$2m_p^2 c^2 + 2m_p E_{lab} = (4m_p^2 + 4m_p m_\pi + m_\pi^2) c^2.$$

Solving for E_{lab} :

$$E_{lab} = \frac{(2m_p^2 + 4m_p m_\pi + m_\pi^2) c^2}{2m_p} = m_p c^2 + 2m_\pi c^2 + \frac{m_\pi^2 c^2}{2m_p}.$$

This exact derivation avoids the complex velocity transformations required by non-covariant methods.


C. Photon Kinematics and Compton Scattering

Massless particles, such as photons, have four-momenta satisfying $P^\mu P_\mu = 0$. Consequently, $E = |p|c$. The interaction of photons with matter provides a clear demonstration of four-vector algebra.

1. Compton Scattering

Compton scattering describes the elastic scattering of a photon by a charged particle (typically an electron). Historically, this provided crucial evidence for the particle nature of light (Compton, 1923).

Figure 1
Compton Scattering Kinematics



Note. The incoming photon $\gamma(K)$ scatters from an electron $e^-(P)$, producing an outgoing photon $\gamma(K')$ at angle θ and a recoiling electron $e^-(P')$.

Let K and P be the initial four-momenta of the photon and electron, and K' and P' be their final momenta. The electron is initially at rest.

$$\begin{aligned} K &= (\hbar\omega/c, k), \\ P &= (mc, 0), \\ K' &= (\hbar\omega'/c, k'), \\ P' &= (E'/c, p'). \end{aligned}$$

Conservation of four-momentum states:

$$K + P = K' + P'.$$

We wish to find the change in photon wavelength as a function of the scattering angle θ . We rearrange the equation to isolate the final electron momentum, which is usually unmeasured:

$$P' = K + P - K'.$$

We square both sides (contracting the four-vectors with themselves):

$$P'^2 = (K + P - K')^2.$$

Expanding the right side and using the invariants $P^2 = P'^2 = m^2 c^2$ and $K^2 = K'^2 = 0$:

$$\begin{aligned}m^2 c^2 &= K^2 + P^2 + K'^2 + 2K \cdot P - 2K \cdot K' - 2P \cdot K'. \\m^2 c^2 &= 0 + m^2 c^2 + 0 + 2K \cdot P - 2K \cdot K' - 2P \cdot K'.\end{aligned}$$

Subtracting $m^2 c^2$ from both sides:

$$K \cdot P - K \cdot K' - P \cdot K' = 0 \Rightarrow P \cdot (K - K') = K \cdot K'.$$

We now evaluate the dot products in the laboratory frame:

$$\begin{aligned}P \cdot K &= mc \frac{\hbar \omega}{c} - 0 \cdot k = m\hbar \omega, \\P \cdot K' &= m\hbar \omega', \\K \cdot K' &= \frac{\hbar \omega}{c} \frac{\hbar \omega'}{c} - k \cdot k' = \frac{\hbar^2 \omega \omega'}{c^2} (1 - \cos \theta).\end{aligned}$$

Substituting these back into the equation:

$$m\hbar(\omega - \omega') = \frac{\hbar^2 \omega \omega'}{c^2} (1 - \cos \theta).$$

Dividing by $m\hbar \omega \omega'$ and using $\omega = 2\pi c/\lambda$:

$$\frac{1}{\omega'} - \frac{1}{\omega} = \frac{\hbar}{mc^2} (1 - \cos \theta).$$

Converting angular frequency to wavelength ($\lambda = 2\pi c/\omega$):

$$\lambda' - \lambda = \frac{h}{mc} (1 - \cos \theta).$$

This is the celebrated Compton shift formula. The quantity h/mc is the Compton wavelength of the electron (2.43×10^{-12} m). This derivation demonstrates the power of the four-vector method: by squaring the momentum conservation equation, we eliminated the relativistic velocity of the recoiling electron entirely, focusing only on the photon variables of interest.

V. INTERACTION DYNAMICS: FEYNMAN DIAGRAMS AND MANDELSTAM VARIABLES

While the conservation of four-momentum provides the fundamental constraints on particle motions, the calculation of interaction probabilities requires a detailed description of the interaction mechanism. This is achieved through Feynman diagrams and the use of Lorentz-invariant kinematic variables known as Mandelstam variables.

A. Feynman Diagrams as Spacetime Trajectories

Feynman diagrams are not merely schematic representations but are rigorous mathematical prescriptions for calculating scattering amplitudes (Feynman, 1949). In these diagrams, standard convention places time on the horizontal axis and space on the vertical axis (or vice versa). Lines represent particles moving through spacetime:

- **External lines** represent real, observable particles entering or leaving the interaction. They satisfy the on-shell condition $P^\mu P_\mu = m^2 c^2$.
- **Internal lines** represent virtual particles (propagators) that exist only transiently during the interaction. For these particles, $P^\mu P_\mu \neq m^2 c^2$.
- **Vertices** are points where particle lines meet, representing the fundamental interaction strength (coupling). Four-momentum is strictly conserved at every vertex.
-

B. Mandelstam Variables

Consider a general two-body scattering process $1 + 2 \rightarrow 3 + 4$, where incoming particles with momenta P_1, P_2 scatter to produce outgoing particles P_3, P_4 . To describe this process in a frame-independent manner, we define three Lorentz-invariant scalars, known as the Mandelstam variables (s, t, u) (Mandelstam, 1958).

1. The s-channel (Center of Mass Energy)

The variable s is the square of the total center-of-mass energy:

$$s = (P_1 + P_2)^2 = (P_3 + P_4)^2.$$

In the Center of Momentum (CM) frame, where $p_1 + p_2 = 0$, this simplifies to $s = (E_1 + E_2)^2/c^2$. A process dominated by s meaning the collision is controlled by the energy available at the CM often involves the creation of a short-lived intermediate resonance that subsequently decays.

2. The t-channel (Momentum Transfer)


The variable t represents the square of the four-momentum transfer between particle 1 and particle 3:

$$t = (P_1 - P_3)^2 = (P_4 - P_2)^2.$$

In elastic scattering, t is related to the scattering angle θ . For small scattering angles (grazing collisions), t is small and negative. The diagram below illustrates a t -channel exchange, typical of Coulomb scattering or Møller scattering.

Figure 2

t-Channel Exchange in Two-Body Scattering



Note. The diagram represents four-momentum transfer q between the two particle lines, with the Mandelstam variable $t = q^2$.

Here, the four-momentum of the virtual photon is $q = P_1 - P_3$. Since the photon is virtual, $q^2 = t < 0$, meaning it is “spacelike.”

3. The u-channel (Crossed Momentum Transfer)

The variable u is defined as:

$$u = (P_1 - P_4)^2 = (P_3 - P_2)^2.$$

This channel becomes relevant when the outgoing particles can be “crossed”. That is, when particle 1 can scatter into the role of particle 4 rather than particle 3. Physically, this arises in processes involving identical particles (such as electron–electron scattering, where one cannot distinguish which outgoing electron corresponds to which incoming electron) or when quantum numbers permit exchange in a crossed configuration. Both the t - and u -channel diagrams must be included in the amplitude, and their interference gives rise to observable effects such as the angular dependence of the Møller scattering cross section.

4. The Sum Rule

These three variables are not independent. For a process involving particles of masses m_1, m_2, m_3, m_4 , they satisfy the identity:

$$s + t + u = \sum_{i=1}^4 m_i^2 c^2.$$

This relation allows experimentalists to fully characterize a scattering event using only two independent variables (usually s and t) and the particle masses.

VI. CONCLUSION

The transition from Galilean relativity to Special Relativity requires a fundamental reimagining of space and time. By treating time as a coordinate on equal footing with space, the four-vector formalism restores the symmetry of physical laws. We have shown that while individual components of energy and momentum vary between observers, the invariant length of four-vectors, such as proper time, rest mass, and the Mandelstam variables, provides a pathway towards calculation as a foundation of objective reality.

From the simple derivation of time dilation to the calculation of particle production thresholds and Compton scattering, four-vectors simplify the complex algebra of high-velocity mechanics. In modern particle physics, where scattering amplitudes are functions of s , t , and u , this formalism is not merely a convenience; it is the essential language of the universe at high energies.

REFERENCES

- Compton, A. H. (1923). A quantum theory of the scattering of X-rays by light elements. *Physical Review*, 21(5), 483–502. <https://doi.org/10.1103/PhysRev.21.483>
- DiSalle, R. (2020). Space and time: Inertial frames. In E. N. Zalta (Ed.), *The Stanford encyclopedia of philosophy* (Winter 2020 ed.). Stanford University. <https://plato.stanford.edu/archives/win2020/entries/spacetime-iframes/>
- Einstein, A. (1905). Zur Elektrodynamik bewegter Körper. *Annalen der Physik*, 322(10), 891–921. <https://doi.org/10.1002/andp.19053221004>
- Feynman, R. P. (1949). Space-time approach to quantum electrodynamics. *Physical Review*, 76(6), 769–789. <https://doi.org/10.1103/PhysRev.76.769>
- Griffiths, D. (2008). *Introduction to elementary particles* (2nd ed.). Wiley-VCH.
- Halzen, F., & Martin, A. D. (1984). *Quarks and leptons: An introductory course in modern particle physics*. John Wiley & Sons.
- Mandelstam, S. (1958). Determination of the pion-nucleon scattering amplitude from dispersion relations and unitarity. General theory. *Physical Review*, 112(4), 1344–1360. <https://doi.org/10.1103/PhysRev.112.1344>
- Navas, S., Amsler, C., Gutsche, T., Hanhart, C., Hernández-Rey, J. J., Lourenço, C., Masoni, A., Mikhasenko, M., Mitchell, R. E., Patrignani, C., Schwanda, C., Spanier, S., Venanzoni, G., Yuan, C. Z., Agashe, K., Aielli, G., Allanach, B. C., Alvarez-Muñiz, J., Antonelli, M., ... Zyla, P. A. (2024). Review of particle physics. *Physical Review D*, 110(3), 030001. <https://doi.org/10.1103/PhysRevD.110.030001>
- Rivadulla, A. (2004). The Newtonian limit of relativity theory and the rationality of theory change. *Synthese*, 141(3), 417–429. <https://doi.org/10.1023/B:SYNT.0000044994.31650.45>
- Rossi, B., & Hall, D. B. (1941). Variation of the rate of decay of mesotrons with momentum. *Physical Review*, 59(3), 223–228. <https://doi.org/10.1103/PhysRev.59.223>
- Tasson, J. (Ed.). (2019). *Symmetry in special and general relativity* [Special issue]. MDPI. https://www.mdpi.com/journal/symmetry/special_issues/Symmetry_Special_General_Relativity
- Taylor, E. F., & Wheeler, J. A. (1992). *Spacetime physics: Introduction to special relativity* (2nd ed.). W. H. Freeman.